

THE FLUORESCENT AND ABSORPTION SPECTRA OF ANTHRACENE AND PHENANTHRENE VAPORS

DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE JOHNS
HOPKINS UNIVERSITY, IN CONFORMITY WITH THE
REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

BY
T. SIDNEY ELSTON

BALTIMORE
1906

THE FLUORESCENT AND ABSORPTION SPECTRA OF ANTHRACENE AND PHENANTHRENE VAPORS

DISSERTATION

SUBMITTED TO THE BOARD OF UNIVERSITY STUDIES OF THE JOHNS
HOPKINS UNIVERSITY, IN CONFORMITY WITH THE
REQUIREMENTS FOR THE DEGREE OF
DOCTOR OF PHILOSOPHY

BY
T. SIDNEY ELSTON

BALTIMORE
1906



Reprinted from
THE ASTROPHYSICAL JOURNAL, Vol. XXV, No. 3
April 1907

PRINTED AT THE UNIVERSITY OF CHICAGO PRESS, CHICAGO

TABLE OF CONTENTS

	PAGE
I. INTRODUCTION	155
II. THE FLUORESCENT SPECTRUM OF ANTHRACENE VAPOR	156
<i>a)</i> Method and Apparatus	
<i>b)</i> The Fluorescence	
III. THE EFFECT OF INCLOSING A FOREIGN GAS WITH THE ANTHRACENE	158
IV. THE EFFECT OF PRESSURE	161
V. THE EFFECT OF TEMPERATURE	161
VI. THE EFFECT OF VARYING THE DENSITY OF THE VAPOR	161
VII. THE ABSORPTION SPECTRUM	162
VIII. THE RELATION BETWEEN THE EXCITING LIGHT AND THE FLUORESCENCE—STOKES'S LAW	163
1. The Carbon Arc as Source	
2. The Zinc Arc and Magnesium Spark as Sources	
3. Stokes's Law	
4. Discussion of Results	
IX. THE FLUORESCENCE AND ABSORPTION OF PHENANTHRENE VAPOR, THE ISOMER OF ANTHRACENE VAPOR	168
X. SUMMARY	168



Digitized by the Internet Archive
in 2026

https://archive.org/details/IA41546428_0044

THE ASTROPHYSICAL JOURNAL

AN INTERNATIONAL REVIEW OF SPECTROSCOPY
AND ASTRONOMICAL PHYSICS

VOLUME XXV

APRIL 1907

NUMBER 3

THE FLUORESCENT AND ABSORPTION SPECTRA OF ANTHRACENE AND PHENANTHRENE VAPORS

By T. SIDNEY ELSTON

I. INTRODUCTION

The fluorescence of vapors was first observed by Lommel¹ for iodine vapor in 1883. A few years later Ramsay and Young² and E. Wiedemann³ found that some liquid solutions, when raised above their critical temperatures, fluoresce. Later, Wiedemann and Schmidt⁴ discovered a long list of fluorescent organic vapors, among which were anthracene and phenanthrene. They found that these vapors under the action of sunlight or light from a carbon arc fluoresced an intense blue, the fluorescent light lying, as a whole, on the red side of the region of maximum absorption. In 1896 they⁵ discovered also that the metallic vapors of sodium and potassium are fluorescent. Quite recently Professor R. W. Wood⁶ has made an extensive study of the fluorescence of sodium vapor. Hartley⁷ has shown that mercury vapor also is fluorescent under certain conditions.

With the exception of sodium vapor, none of these vapors had been investigated extensively for its fluorescence; and yet it seemed

¹ *Wied. Ann.*, **19**, 356, 1883.

² *Chem. News*, **54**, 203, 1886.

³ *Wied. Ann.*, **41**, 209, 1890.

⁴ *Ibid.*, **56**, 18, 1895.

⁵ *Ibid.*, **57**, 447, 1896.

⁶ *Phil. Mag.*, **10**, 513, 1905.

⁷ *Proc. R. S.*, **76** A, 428, 1905.

desirable that such work should be done, for the sake of the aid which it might afford for the solution of the problem of luminescence, and ultimately of the problem of the nature of matter. In the present investigation the primary object was to study in more detail the fluorescence of anthracene vapor, one of the very strongly fluorescent organic vapors. Later a similar study was made of phenanthrene because of the close resemblance between its fluorescent spectrum and that of anthracene, its isomer.

II. THE FLUORESCENT SPECTRUM OF ANTHRACENE VAPOR

a) *Method and apparatus.*—The fluorescence of the anthracene vapor was first studied by means of a small quartz spectrograph. The pure anthracene used was obtained from the commercial product by distilling twice from an excess of caustic potash, and then crystallizing from benzene under the action of the sunlight. A few centigrams of the crystals were inclosed in a crown-glass bulb (5 to 8 cm in diameter), which was then evacuated and sealed.

The bulb was placed in a heating apparatus arranged as shown in Fig. 1. It was hung in a wire cage between the two side tubes, T_1 and T_2 , in such a manner that the light employed to excite fluorescence, after passage through the condensing lens, L , and reflection from the mirror, M , downward through the mica window, W , was focused at the center of the bulb.

The tube, T_1 , contained a rectangular opening at its inner end and a glass window near its outer end. When the bulb was heated and the anthracene made to fluoresce, the fluorescent light was examined by the spectrograph through this tube.

The larger tube, T_2 , provided with a cap over its outer end, served a double rôle: first, as an observing tube for adjusting the apparatus so as properly to focus the light in the bulb; and, secondly, as a dark background against which to observe the fluorescence through the other tube, T_1 .

Care was always taken to adjust the bulb so that the cone of exciting light should be reflected from the bottom of the bulb straight back along its path. This was done in order to get rid, as far as possible, of the light scattered by reflection from the sides of the bulb. It was found, when bulbs of the proper size and shape were adjusted

in this way, that an exposure of three hours, made for scattered light alone, gave a very faint photograph as compared with that of the fluorescent and scattered light combined.

Various sources of violet and ultra-violet light were tried—viz., the sun, the carbon and zinc arcs in air, the mercury, silver, and lead arcs *in vacuo*, and the cadmium, aluminium, copper, zinc, and magnesium sparks. Of these the carbon arc produced the most




FIG. 1

intense fluorescence in the anthracene vapor, and so was employed almost exclusively. It was found that the fluorescence was due more to the light from the incandescent gases between the carbons than to the incandescent carbons themselves. Its intensity increased or decreased with the intensity of the group of three cyanogen bands in the violet.

b) *The fluorescence.*—When the bulb was heated until the anthracene began to vaporize, the path of the arc-light through it was

marked by a cone of illuminated particles which scattered partially polarized white light in all directions. Gradually this disappeared as the anthracene became completely vaporized, and was replaced by a cone of brilliant blue fluorescent light in which no polarization could be detected. Examined photographically with the quartz spectrograph, it gave a spectrum (Plate X, 1a) which extended continuously from $365\text{ }\mu\mu$ to $470\text{ }\mu\mu$, with its regions of maximum intensity located at 390 , 415 , and $432\text{ }\mu\mu$. The same spectrum was also obtained when sunlight, or the light from any of the other sources was employed as the exciting light.

The range of the apparatus was from $\lambda\text{ }325$ to $540\text{ }\mu\mu$, the limit in the ultra-violet being set by the crown glass of the bulb which is opaque to radiation of wave-length shorter than $325\text{ }\mu\mu$.


The photograph did not show the slightest evidence of lines in the fluorescent spectrum. This, it was thought, might be due to the small resolving power of the quartz spectrograph in the region of the fluorescence; so a photograph was taken with a special three-crown-prism spectrograph, designed by Professor Wood for his work on sodium vapor. The resolving power of this instrument was about fifteen times that of the quartz spectrograph; but in this case also the photograph failed to show any lines in the spectrum. The fluorescence was not intense enough to warrant the use of instruments of higher resolving power. It seems highly probable, however, that the fluorescent spectrum of anthracene vapor is not resolvable into lines like that of sodium vapor, but resembles, instead, the banded spectra of fluorescent liquids and solutions.

III. THE EFFECT OF INCLOSING A FOREIGN GAS WITH THE ANTHRACENE

To determine what effect upon the fluorescence the presence of another gas would have, the anthracene was inclosed in the bulb in turn with various gases at atmospheric pressure. It was found that some gases had practically no effect, while others extinguished the fluorescence more or less completely. The results are given in the accompanying table.

In the case of the gases listed in the first column of the table the quality of the fluorescence was unaffected by their introduction into

PLATE X



FLUORESCENT AND ABSORPTION SPECTRA OF ANTHRACENE AND PHENANTHRENE VAPORS

1. *a*, Fluorescence of Anthracene; *b*, Diffused Light; *c*, Cadmium Spark.
2. *a*, Nernst Filament; *b*, Absorption of Anthracene; *c*, Cadmium Spark.
3. *a*, Nernst Filament; *b*, Absorption of Anthracene (dense vapor); *c*, Cadmium Spark.
4. *a*, Nernst Filament; *b*, Absorption, Commercial Anthracene; *c*, Cadmium Spark.
5. *a*, *g*, Cadmium Spark; *h*, Carbon Arc Spectrum; *d*, Solution A as screen; *e*, Solution B; *f*, Solution C; *g*, Crown-Glass Plate as screen.
6. *a*, Fluorescence of Phenanthrene; *b*, Cadmium Spark.

FLUORESCENCE OF ANTHRACENE VAPOR

Not Affected by	Extinguished by	Weakened by
Nitrogen Hydrogen Illuminating gas Carbon monoxide Carbon dioxide Mercury vapor	Cyanogen Chlorine Sulphur dioxide Oxygen	Air

the bulb. A slight decrease in intensity, however, was observed, due, probably, to the higher pressures in the bulb after the gases were introduced (see § IV). So far as could be ascertained, none of these gases reacts chemically upon the anthracene vapor within the range of temperatures set by the apparatus.

In the case of the gases listed in the second column the effect was practically to extinguish the fluorescence. Only the faintest trace, if any, of fluorescence could be observed when they were mixed with the anthracene. For these gases it was found that chemical action began either as soon as the temperature was raised to a point where the anthracene was vaporized (this was the case with cyanogen and chlorine), or at a temperature not much higher (this was the case with sulphur dioxide and oxygen). It seemed highly probable from this that the extinction of the fluorescence was due to some sort of chemical influence.

That the chemical influence, if such it be, which affected the fluorescence, need not be such as to produce a permanent chemical change in the anthracene was shown in the following way: A bulb containing anthracene and oxygen was placed in an air-bath with a similar bulb containing only anthracene. The temperature was then raised just high enough to obtain strong fluorescence in the second bulb, but not high enough to cause any visible chemical reaction between the oxygen and the anthracene in the first bulb. The first bulb was tested for fluorescence and found not to show the slightest traces. It was then removed and the oxygen pumped out, the crystallized anthracene being left behind, and the bulb again placed in the bath and tested. The fluorescence in this bulb was now found to be as brilliant as in the other one. To find out if some of the anthracene had not been acted upon and a consequent change in

volume of the oxygen produced, a bulb was used which terminated in a long capillary tube. The oxygen gas and anthracene crystals were inclosed at atmospheric pressure and the bulb sealed. Then, after subjecting the bulb to the same heating as before, the end of the capillary tube was broken off under mercury, and the oxygen again brought to atmospheric pressure. Not the slightest change in its volume was observed. If any chemical action had occurred, it was evidently of such a nature as to produce compensating changes in the volume of the reacting substances. As this is highly improbable, we are led to conclude that no reaction took place between the anthracene and the oxygen, but that whatever mutual action occurred produced effects which lasted only while the given conditions of temperature and mutual contact were maintained. During this time practically all of the anthracene molecules must have been affected, for the extinction of the fluorescence was complete throughout the vapor.

The foregoing facts may be explained by assuming that, when anthracene vapor is mixed with oxygen at a temperature not much below that at which the two react chemically, there is a preliminary grouping of the oxygen molecules about the anthracene molecules, or vice versa, and that, while thus associated and before they rush together into closer contact to form a new chemical compound, the mutual forces are such as to prevent the anthracene molecules from fluorescing, but are not great enough to bring about a reaction. As thus considered, the phenomenon is simply the first stage of a chemical reaction.

The effect of oxygen upon the anthracene vapor is in many ways similar to that of many liquid solvents upon the fluorescent substances dissolved in them.¹

When the vapor was mixed with air, the fluorescence was much weakened, becoming almost too weak to be seen when air at atmospheric pressure or greater was used. This weakening was easily traceable to the oxygen contained in the air.

It seems highly probable from the foregoing facts that the presence of foreign molecules, heavy or light, among the fluorescing anthracene molecules does not have any appreciable effect upon the

¹ Kauffmann and Beiswenger, *Zeitschrift für phys. Chem.*, 50, 350, 1904.

fluorescence at ordinary pressures, except in those cases where a chemical change, either incipient or permanent, occurs.

IV. THE EFFECT OF PRESSURE

In the experiment with the illuminating gas a thick-walled glass bulb was employed, and the pressure was varied from 1 to 12 atmospheres. The visible fluorescence became less and less intense as the pressure was increased, but was still perceptible at the highest pressure. So far as this was tried with the other gases, the same results were obtained, an increase of pressure in each case causing a diminution in the intensity of the fluorescence. The change in intensity was not very marked until the pressure exceeded 1 atmosphere. At all pressures below this the intensity of the fluorescence was practically independent of the pressure.

The quality of the fluorescence was not affected by the pressure; the same spectrum was obtained at high as at low pressures.

V. THE EFFECT OF TEMPERATURE

The temperature of the bulb was varied from the temperature at which the vapor began to fluoresce (the boiling-point of anthracene is 351°C.) to the temperature at which the glass began to soften (the melting-point of the glass bulb is about 1000°C.). As the temperature was raised the intensity of the fluorescence diminished. This was the case whether the bulb contained the pure anthracene alone or anthracene mixed with an inert gas. Of course, as the temperature was raised, the pressure of the inclosed vapor was increased; so the effect may have been due to this.

VI. THE EFFECT OF VARYING THE DENSITY OF THE VAPOR

When the amount of anthracene inclosed in the bulb was gradually increased, and the vapor density thus correspondingly increased, it was found that the cone of visible fluorescence, which at first extended completely through the bulb from the point where the exciting light entered to the point where it left, was foreshortened in the direction of its length, until finally it was shrunk down almost to the surface where the exciting light entered the bulb. At the same time the intrinsic brightness of the fluorescence was diminished, due, undoubtedly, to the consequent increase in pressure.

The foreshortening of the cone of fluorescence means, of course, that the particular radiation which excites the visible fluorescence is completely absorbed out of the incident light before this penetrates very far into a bulb filled with dense anthracene vapor.

The quality of the fluorescence was not affected, as the spectrum photographs for bulbs containing different amounts of anthracene showed.

VII. THE ABSORPTION SPECTRUM

To obtain the absorption spectrum of the vapor, it was found convenient to use a long glass tube in which to inclose the anthracene. This was placed inside a cylindrical iron tube with a glass window at each end, as shown in Fig. 2.




FIG. 2

The light from a Nernst filament was focused in the tube at one end, and, on emerging at the other end, was examined photographically with the quartz spectrograph.

The photograph (Plate X, 3*b*) was taken when the anthracene vapor in the tube was quite dense. It shows that the pure dense vapor absorbs continuously from λ 425 to 325 $\mu\mu$, the limit set by the crown glass. When the tubes containing less anthracene, or the bulbs employed in getting fluorescence, were used, the absorption did not extend so far down toward longer wave-lengths (Plate X, 2*b*).

The absorption of the vapor was also studied with a 12.5-foot concave Rowland grating, but with the same result. There was no evidence of lines in either case; the absorption consisted of a continuous region between λ 325 $\mu\mu$ and about 425 $\mu\mu$.

Purified anthracene was used in obtaining the absorption spectra just described. When the commercial anthracene (Eimer & Amend, 90 per cent.) was used, the absorption spectrum obtained (Plate X,

4b) was found to be the same as for the pure anthracene, with the exception that there was a narrow band at wave-length $452\ \mu\mu$.

VIII. THE RELATION BETWEEN THE EXCITING LIGHT AND THE FLUORESCENCE—STOKES'S LAW

1. *The carbon arc as source.*—In the earlier stages of the investigation (when the fluorescent spectrum was excited by the light from a carbon-arc lamp and the photograph taken showed that it consisted of three bands which were somewhat similarly spaced, but farther toward the red, as compared with the three intense cyanogen bands of the arc in that region of the spectrum), it was thought probable that the three cyanogen bands had each excited its own fluorescent band in the anthracene vapor, thus giving rise to the three bands found in the fluorescent spectrum. The fact, that sunlight excited the same bands seemed to make this doubtful; but it was not until the work, now to be described, was done that it was shown that, in all probability, the entire fluorescence of the vapor excited by light from the carbon arc was largely due to the strong lines in the head of the middle one of the three cyanogen bands, the one whose head lies at wave-length $388.3\ \mu\mu$.

Two methods were employed in determining the approximate wave-length of the exciting light.

a) The first to be tried was what might be called the "method of exclusion." It consisted in screening off different portions of the spectrum in turn, by means of properly chosen absorbing solutions, and observing the effect upon the fluorescence. By referring to Dr. Uhler's photographs of the absorption spectra of various solutions, the following solutions were chosen and found to answer the purpose:

A. Nitroso-dimethyl-aniline solution, 2 parts by volume of saturated solution to 3 parts of water.

B. Potassium permanganate solution, 1 part by volume of saturated solution to 50 parts of water.

C. Naphthol disulfonic acid solution, 2 parts by volume of saturated solution to 5 parts of water.

These solutions were introduced into a glass cell (1 cm thick) which was placed in the path of the exciting light between the condensing lens (see Fig. 1) and the bulb.

When solution A was employed as an absorbing screen, the anthracene vapor failed to fluoresce. Now, this solution has a strong absorption band extending from 375 to 480 $\mu\mu$ (see Plate X, 5*d*), while it transmits quite freely on both sides of the band, within the limits set by the glass. It is evident from this that the light which causes the fluorescence in the vapor is located somewhere in the region 375–480 $\mu\mu$. It is in this region that two of the cyanogen bands are located; the third lies just outside of it toward the ultra-violet.

The solution B is transparent to light of wave-length 395–465 $\mu\mu$, but is quite opaque to the rest of the spectrum in the given region. When this solution was substituted in the cell for solution A, no fluorescence could be observed. This shows that the light producing the fluorescence does not lie between 395 and 465 $\mu\mu$, and therefore must lie either in the region 465–480 $\mu\mu$, or in the region 375–395 $\mu\mu$, or in both. That it does not lie in the region 465–480 $\mu\mu$ is probable from the fact that the carbon arc spectrum is relatively weak in this region. We are then led to conclude that the exciting light lies somewhere between the wave-lengths 375 and 395 $\mu\mu$. It is in this region that the strong lines in the middle one of the three cyanogen bands are located.

b) A confirmation of the foregoing results was obtained in the following way: A slit, two condensing lenses, and a large crown-glass prism were arranged so as to give a comparatively pure and bright spectrum of the arc. A slitted screen, with the bulb containing the anthracene vapor behind it, was then passed from point to point along through the spectrum. Starting at the red end there was no visible evidence of fluorescence in the vapor until the bulb was brought into that part of the spectrum corresponding to wave-lengths 390–375 $\mu\mu$, when it began to fluoresce with the characteristic brilliant blue color, the cone of fluorescence extending entirely through the bulb. When moved on into the region of the cyanogen band whose head is at 359 $\mu\mu$, a dim fluorescence was observed which penetrated only a few millimeters into the bulb. This band had played small part, therefore, in the photographs of the fluorescent spectrum previously taken, where the collimator of the spectrograph had been directed toward the center of the bulb.

The slitted screen was now adjusted so as to allow only the light

near the head of the cyanogen band at $388.3\ \mu\mu$ to enter the bulb. A photograph of the fluorescence produced was taken which gave the same bright bands and continuous regions shown in the former photographs (see Plate X, 1a).

The slit was then moved so as to admit to the bulb the head of the cyanogen band at $359\ \mu\mu$, and a photograph of the fluorescence was taken. It showed that the spectrum was the same as that excited by the other cyanogen band.

The relative spectral positions of the fluorescence and the exciting lights for the two experiments just described are shown in Fig. 3. The block E_1 represents the spectral width and location of the exciting light in the first experiment, and F_1 the general form of distribution of intensity in the fluorescence which it excited in the anthracene vapor, while E_2 and F_2 represent the corresponding quantities in the second experiment. The two curves have been deduced from the photographs and indicate that the fluorescence in the two cases was the same in every way except in absolute intensity.

2. *The zinc arc and magnesium spark as sources.*—To obtain further evidence of the relation between the fluorescence and the wave-length of the light which excites it, two sources were employed which gave regions of exciting light quite widely separated. The first was the zinc arc, which gives an isolated group of very intense lines in the region $328\text{--}335\ \mu\mu$, just within the ultra-violet limit set by the absorption of the crown-glass bulb; the second was the magnesium spark, which gives an intense isolated triplet in the region $383\text{--}385\ \mu\mu$, not very far from the limit toward the red beyond which the light ceases to have power to excite fluorescence in the anthracene vapor. This limit was earlier found to be in the neighborhood of wave-length $400\ \mu\mu$.

Between 325 and $400\ \mu\mu$ the arc-spectrum of zinc contains, besides the group just mentioned, only three very faint lines whose aggregate intensity is less than 1 per cent. of that of the group. If the zinc arc is used, therefore, to excite the fluorescence in the anthracene, the exciting light is practically all located in the region $328\text{--}335\ \mu\mu$. When this was done (the zinc arc being substituted for the carbon arc) and the fluorescence produced in the bulb was photographed, it was found to give the same fluorescent spectrum as the carbon

arc. The block E_3 and the curve F_3 in Fig. 3 represent respectively the exciting light and the fluorescence for this case.

In the case of the magnesium spark there are a few faint lines in the region 325-400 $\mu\mu$ besides the triplet mentioned, but their aggre-




FIG. 3

gate intensity is negligible in comparison with that of the triplet, so that practically all of the exciting light from a pure magnesium spark is in the region 383-385 $\mu\mu$. It was found, however, that the elec-

trodes used contained zinc as an impurity. In order to screen off the lines due to the zinc, the solution C (see § VIII, 1) was used as an absorbing screen. It is opaque to ultra-violet light from 360 on out beyond 325 $\mu\mu$, and therefore to the group of zinc lines at 328–335 $\mu\mu$. When the magnesium spark, as thus screened, was used as the source and the fluorescence excited was photographed, it was found to give the same spectrum as the carbon and zinc arcs. The block E_4 and the curve F_4 in Fig. 3 represent respectively the exciting light and the fluorescence for this case.

In connection with the results given above, it should be remembered that the anthracene vapor used absorbs continuously from about 400 $\mu\mu$ to some point beyond 325 $\mu\mu$.

3. *Stokes's law*.—It is evident from the foregoing facts that Stokes's law, which states that the fluorescent light is of longer wave-length than the light which excites it, is not strictly true in the case of anthracene vapor; for in at least two cases (1 and 4 in Fig. 3) part of the fluorescent light is of shorter wave-length than the corresponding exciting light. In a general sense, however, the law may be considered as fairly representing the facts, since the fluorescent spectrum as a *whole* is of longer wave-length than the exciting light.


4. *Discussion of results*.—We are led to conclude from the foregoing facts, (a) that the fluorescence of anthracene vapor is excited by light situated anywhere in the ultra-violet region of absorption of the vapor, and (b) that the character of the fluorescence is entirely independent of the source and wave-length of the light which excites it.

If we assume that the fluorescence is produced by a system of electrons within the molecule, then, in order to account for the fluorescent spectrum of anthracene vapor, which, as we found, was composed of three bands superposed upon a continuous spectrum, we may consider either that the electrons corresponding in period to the three bands are more numerous than those which give rise to the weaker, continuous parts of the fluorescence, or that the former are set in more violent vibration; also, that the system of electrons is so intimately connected in its parts that, when disturbed by the exciting light in any manner, be it direct or through an intermediary "luminophore," *all* the electrons in the system are set in vibration. If the disturbance of the system takes place through an intermediary

"luminophore,"¹ as seems more probable, then this luminophore undoubtedly consists of a connected system of electrons whose periods correspond to those of the absorption spectrum of the vapor.

IX. THE FLUORESCENCE AND ABSORPTION OF PHENANTHRENE VAPOR, THE ISOMER OF ANTHRACENE VAPOR

Phenanthrene is a substance which has the same chemical composition, $C_{14}H_{10}$, as anthracene, but a slightly different structural composition as the following figure shows:



The only difference in structure, according to this, is in the manner in which the four groups or radicals are linked together.

When the fluorescent spectrum of pure phenanthrene vapor was photographed (Plate X, 6a), it was found to consist of the same bands as that of anthracene, but with an additional band at $360\ \mu\mu$.

Its absorption spectrum was photographed and found to be the same as that of the anthracene vapor.

It would seem from these facts that there is an intimate connection between the fluorescence of the vapors of the two substances, undoubtedly due to their common chemical composition and similar structural composition. Just what gives rise to the extra band in the fluorescent spectrum of phenanthrene vapor is not apparent.

X. SUMMARY

The following is a summary of the results of this investigation:

(1) The fluorescent spectrum of anthracene vapor consists of three bright bands at $\lambda\ 390, 415$, and $432\ \mu\mu$ superposed upon a continuous region extending from 365 to $470\ \mu\mu$. There is no evidence of lines. The absorption spectrum extends continuously from about $400\ \mu\mu$ to some point beyond $325\ \mu\mu$.

(2) The presence of inert gases in the vapor does not affect the fluorescence, so long as the pressure is below an atmosphere. But

¹ W. Kauffmann, *Ber. d. Phys. Gesell.*, **21**, 375, 1905.

such gases as oxygen, chlorine, and sulphur dioxide, which at high temperatures react chemically with anthracene, almost completely extinguish the fluorescence at ordinary pressures.

(3) The intensity of the visible fluorescence decreases as the pressure of the gas inclosed with the anthracene is increased. It is not marked, however, at pressures lower than an atmosphere. The quality of the fluorescence is not affected by a change in pressure.

(4) Increasing the density of the fluorescing vapor has no effect upon the quality of the fluorescence, but diminishes its intensity slightly.

(5) The fluorescence of anthracene vapor may be excited by light located anywhere within the ultra-violet region of absorption of the vapor.

(6) The same fluorescent spectrum is produced independent of what the source or wave-length of the exciting light may be, provided the condition noted in (5) is observed.

(7) There is an intimate connection between the fluorescence of anthracene vapor and its isomer, phenanthrene vapor.

This investigation was begun at the suggestion and has been carried out under the direction of Professor R. W. Wood, to whom I wish to extend my thanks for the interest which he has shown, and the encouragement and help which he has given me in the course of the work. I wish also to express my gratitude to Professor J. S. Ames for his unfailing courtesy and kindness.

JOHNS HOPKINS UNIVERSITY

BIOGRAPHY

Thomas Sidney Elston was born in the town of Woodland, Sacramento Valley, Cal., on the sixteenth of December, 1872. He received his preparatory education at Hesperian College, Woodland, Cal. Between the date of his graduation from this school and his matriculation in 1895 at the University of California at Berkeley he spent three years in an accountant's office. He received his bachelor's degree from the Colleges of Natural Science at the university in 1899. The following year he taught physics in the San Diego High School, California. The second year he was called to take charge of the same work in the high school at Berkeley, in which he continued until he came to the Johns Hopkins University as Research Assistant to Professor R. W. Wood. Since that time he has been pursuing graduate study in Physics in the Johns Hopkins Physical Laboratory under Professors Ames and Wood. His first and second minors have been respectively Mathematics under Dr. Cohen and Physical Chemistry under Dr. Jones.

